

ALGORITHM FOR GENERALIZED GARMAN EQUATION IN OPTION PRICING OF A FINANCIAL DERIVATIVES WITH STOCHASTIC VOLATILITY MODELS

Socaciu Tiberiu

University Stefan cel Mare of Suceava Faculty of Economics and Public Administration, Computer Science Chair Suceava, str. Universitatii nr. 13, Corp H

and

West University Vasile Goldis of Arad Computer Science Faculty, Computer Science Chair Arad, Bd. Revolutiei nr. 81 e-mail: socaciu@seap.usv.ro tel: 0744 820 827

Danubianu Mirela

University Stefan cel Mare of Suceava Faculty of Electrical Engineer and Computer Science, Computers Chair Suceava, str. Universitatii nr. 13, Corp C e-mail: mdanub@eed.usv.ro tel: 0744 820 827

Maxim Ioan

University Stefan cel Mare of Suceava Teachers' Training Department Suceava, str. Universitatii nr. 13, Corp A e-mail: maximioan@yahoo.com tel: 0744 820 827

Naaji Antoanela

West University Vasile Goldis of Arad Computer Science Faculty, Computer Science Chair Arad, Bd. Revolutiei nr. 81 e-mail: anaaji@uvvg.ro tel: 0744 820 827

In our paper we build a recurrence from generalized Garman equation and discretization of 3-dimensional domain. From recurrence we build an algorithm for computing values of an option based on time, momentan volatility of support and value of support on a spot market of an exchange.

Keywords: financial derivatives, Black-Scholes PDE, Garman PDE, recurrence, algorithm

JEL classification: G13, C69

1. Garman functions attached on generalized model with stochastic volatility

In Black-Scholes model⁶²³ we suppose that volatility is a constant value. Unfortunately, volatility is not constant, and more, can not be predicted for periods greater than several months or even can not be observed directly. For these reasons, it seems natural to present itself as the volatility of a random variable that satisfies a stochastic process. Assume that the variation rate asset support satisfy a stochastic differential equation (SDE) of the form:

$$dS(t) = A(t, S(t), V(t))dt + B(t, S(t), V(t))dW_1(t) \quad (1)$$

but in addition we assume that the square of volatility (v), follows the stochastic process:

$$dv(t) = C(t, S(t), V(t))dt + D(t, S(t), V(t))dW_2(t), \quad (2)$$

where the two *Wiener processes* $W_1(t)_{t \geq 0}$ și $W_2(t)_{t \geq 0}$ are correlated:

$$(dW_1(t))(dW_2(t)) = \rho dt.$$

In this case the value of an option is a function of three variables, $V(t, S, v)$ and because we have two sources of risk need to build a portfolio of coverage to include the option in question and two other components. Assume the existence of a derivative financial unca on the same support, so that the value of this option is a function of three variables, $V_1(t, s, v)$ to check with a law counterparts of $V(t, S, v)$. After applying extended lemma of Ito⁶²⁴, we obtain the existance of a special value β named as *the market price of volatility risk*⁶²⁵, where:

$$\beta = (-rV + V_s rS + V_t + AV_s + CV_v + \frac{1}{2}B^2 V_{SS} + \frac{1}{2}D^2 V_{vv} + \rho BDV_{sv} - AV_s) / (DV_v) \quad (3)$$

If β is known, (3) rewrite as:

$$V_t + rSV_s + (C - D\beta)V_v + \frac{1}{2}B^2 V_{SS} + \frac{1}{2}D^2 V_{vv} + \rho BDV_{sv} - rV = 0 \quad (4)$$

623 see [Black1973].

624 see [Socaciu2009a].

625 see [Moodley2005], p. 8.

is known as *Garman equation*. Note that (4) is a partial differential equation (PDE), like *Black-Scholes* equation, obtained from SDE of model.

2. Particular cases of Garman functions

1. *Black-Scholes*; in conditions:

$$\begin{aligned} B^2 &= vS^2 \\ C = D &= 0 \Leftrightarrow dv = 0 \text{ (constant volatility)} \\ v &= \sigma^2 \end{aligned}$$

we re-obtain Black-Scholes equation⁶²⁶:

$$V_t + rSV_S + \frac{1}{2}\sigma^2 S^2 V_{SS} - rV = 0$$

2. *Heston*; in conditions:

$$\begin{aligned} A &= \mu S \\ B &= v^{1/2} S \\ C &= k(\theta - v) \\ D &= \xi v^{1/2} \\ \beta &= \lambda v^{1/2} \end{aligned}$$

we re-obtain⁶²⁷:

$$V_t + rSV_S + (k(\theta - v) - \xi \lambda v) V_v + \frac{1}{2}vS^2 V_{SS} + \frac{1}{2}\xi^2 v V_{vv} + \rho S \xi v V_{Sv} - rV = 0 \quad (5)$$

3. *other cases* for particular models can be obtained via particular A, B, C, D and β . An incomplete list of models⁶²⁸ can be retrieved from [Sundaresan2000]⁶²⁹.

3. Garman equation on domains' frontier

Equation (5) has on frontiers⁶³⁰:

$$V(t, 0, v) = 0, \text{ with } 0 \leq t \leq T, \text{ all } v. \quad (6)$$

and:

$$V(T, S, v) = \text{payoff}(S), \text{ with } 0 \leq S, \text{ all } v. \quad (7)$$

We need add a new condition on v axis. Simplest condition is based on zero-volatility. In this case, (2) is vanished, and (1) will be rewrite as a regular BS equation with a specific solutions like⁶³¹:

$$V(t, S, 0) = \text{BS}(t, S) \quad (8)$$

4. Domain's discretization and recurrence obtained from Garman equation

We will apply a *Finite Element Method* (FEM⁶³²) for resolve (5). We know that FEMs are main method for resolve PDE⁶³³. First step is to reduce initial domain D_i :

$$D_i = [0, T] \times \mathbb{R}_+ \times \mathbb{R}_+$$

626 see [Black1973].

627 see [Lazăr2006], [Mikhailov2003].

628 with specification of A and B.

629 an other incomplete list can be retrieved in [Lazăr2006].

630 like in [Socaciu2009b].

631 Numerical solutions will be presented in [Socaciu2009a], [Socaciu2009b] and [Socaciu2009c].

632 see [Gârbea1990].

633 see [Coman1994].

to a *limited domain* D:

$$D = [0, T] \times [0, \text{MaxS}] \times [0, \text{MaxV}],$$

where MaxS is a *big number* that cover all values of potential quotations, and MaxV is an big number that cover all potential volatilities. After discretization with Δt , ΔS and Δv steps we will obtain a grid of points. We denote:

$$V[i,j,k] = V(i(\Delta t), j(\Delta S), k(\Delta v)), \text{ with } 0 \leq i \leq M, 0 \leq j \leq N, 0 \leq k \leq P$$

where number of points on axis:

$$\begin{aligned} M &:= \text{MaxS}/(\Delta S) \\ N &:= T/(\Delta t) \\ P &:= \text{MaxV}/(\Delta v) \end{aligned}$$

than (4) will be rewrite as:

$$\begin{aligned} &(V[i,j,k]-V[i-1,j,k])/(\Delta t)+rj(\Delta S)(V[i,j+1,k]-V[i,j-1,k])/(2(\Delta S)) \\ &+(C-D\beta)(V[i,j,k+1]-V[i,j,k-1])/(2(\Delta v)) \\ &+1/2B^2(V[i,j+1,k]-2V[i,j,k]+V[i,j-1,k])/(\Delta S)^2 \\ &+1/2D^2(V[i,j,k+1]-2V[i,j,k]+V[i,j,k-1])/(\Delta v)^2 \\ &+\rho BD(V[i,j+1,k+1]-V[i,j+1,k-1]-V[i,j-1,k+1]+V[i,j+1,k+1])/((\Delta S)(\Delta v)) \\ &-rV[i,j,k] = 0 \end{aligned}$$

That can be rewrite as:

$$\begin{aligned} &a(V[i,j,k]-V[i-1,j,k])+b(V[i,j+1,k]-V[i,j-1,k])+c(V[i,j,k+1]-V[i,j,k-1]) \\ &+d(V[i,j+1,k]-2V[i,j,k]+V[i,j-1,k])+e(V[i,j,k+1]-2V[i,j,k]+V[i,j,k-1]) \\ &+f(V[i,j+1,k+1]-V[i,j+1,k-1]-V[i,j-1,k+1]+V[i,j+1,k+1])-gV[i,j,k] = 0 \end{aligned} \quad (9)$$

where:

$$\begin{aligned} a &= (\Delta t)(\Delta S)^2(\Delta v)^2 \\ b &= 1/2rj(\Delta t)(\Delta S)^2(\Delta v)^2 \\ c &= 1/2(C-D\beta)(\Delta t)(\Delta S)^2(\Delta v) \\ d &= 1/2B^2(\Delta t)(\Delta v)^2 \\ e &= 1/2D^2(\Delta t)(\Delta v)^2 \\ f &= \rho BD(\Delta t)(\Delta v)(\Delta S) \\ g &= r(\Delta t)(\Delta v)^2(\Delta S)^2 \end{aligned}$$

But (9) can be rewrite in explicit form:

$$\begin{aligned} V[i-1,j,k] &= pV[i,j,k]+qV[i,j+1,k]+rV[i,j-1,k]+sV[i,j,k+1]+tV[i,j,k-1] \\ &+uV[i,j+1,k+1]-uV[i,j+1,k-1]-uV[i,j-1,k+1]+uV[i,j+1,k+1] \end{aligned} \quad (10)$$

where:

$$\begin{aligned} p &= (a-2d-2e-g)/a \\ q &= (b+d)/a \\ r &= (d-b)/a \\ s &= (c+e) \\ t &= (e-c) \\ u &= f/a \end{aligned}$$

Note that coefficients p-u are based on t, S and v, and we must rewrite all of them in spirit of (i,j,k).

5. Numerical algorithm based for Garman equation based on triple recurrence

Numerical algorithm is based on (10) recurrence with supplementally (6)-(8) frontier condition is:

```
For i:=MaxM downto 0 do
  For j:=0 to MaxN+MaxM do
    For k:=0 to MaxP+MaxM do
      If i=MaxM then
        V[i,j,k]:=payoff(j(ΔS))
      ElseIf j=0 then
        V[i,j,k]:=0
      ElseIf k=0 then
        V[i,j,k]:=BS(i,j)
      Else
        V[i,j,k]:=pV[i+1,j,k]+qV[i+1,j+1,k]
        +rV[i+1,j-1,k]+sV[i+1,j,k+1]+tV[i+1,j,k-1]
        +uV[i+1,j+1,k+1]-uV[i+1,j+1,k-1]-uV[i+1,j-1,k+1]
        +uV[i+1,j+1,k+1]
      End If
    End For
  End For
End For
```

where BS is a function that call a numerical solution of Black-Scholes like equation on zero-volatility condition. Note that iteration will be computed for j to MaxN+MaxM and for k to MaxP+MaxM because we have some “out of range” problems if we compute $V[0, \text{MaxN}, \text{MaxP}]$ and cheap resolution is to compute a supradomain of D.

6. Further works

We work on obtaining parallel version of this algorithm with slicing of D like in [Socaciu2009a], [Socaciu2009b] and [Socaciu2009c]. We try to obtain similar formulae for a multi-SDE model, like Chen⁶³⁴. We want to start a parallel project for obtain on symbolic way an extended Ito type lemma⁶³⁵ and from this lemma to build an Garman-type equation.

References

- [Black1973] Fischer Black, Myron Scholes, The Pricing of Options and Corporate Liabilities, in Journal of Political Economy, 81 (3), pp. 637–654.
- [Chen1996] Lin Chen, Stochastic Mean and Stochastic Volatility -- A Three-Factor Model of the Term Structure of Interest Rates and Its Application to the Pricing of Interest Rate Derivatives, Blackwell Publishers, 1996.
- [Coman1994] Gheorghe Coman, Analiza numerică, Ed. Libris, Cluj, 1994.
- [Gârbea1990] Dan Gârbea, Analiza cu elemente finite, Editura Tehnica, Bucuresti, 1990, 248 pagini.
- [Iftimie2008] Bogdan Iftimie, Financial models in continuous time, Editura MatrixRom, Bucuresti, 2008, ISBN 978-973-755-350-8, 126 pagini.
- [Lazăr2006] Lazăr Vasile Lucian, Modele de evaluare a opțiunilor, in Studia universitatis Stiinte Economice, Nr. (5) an 2006, volumul II, online at
- http://www.uvvg.ro/studia/economice/index.php?categoryid=10&p2_articleid=74&p142_dis=3&p142_template=Default.
- [Mikhailov2003] Mikhailov, S., Nogel, U., Heston's stochastic volatility model. Implementation, calibration and some extensions, in Wilmott, 2003, 74–94, online at
- http://www.wilmott.com/pdfs/051111_mikh.pdf.
- [Moodley2005] Nimalin Moodley, The Heston Model: A Practical Approach with Matlab Code, University of the Witwatersrand, Johannesburg, South Africa, online at <http://www.math.nyu.edu/~atm262/fall06/compmethods/a1/nimalinmoodley.pdf>.
- [Sundaresan2000] Suresh M. Sundaresan, Continuous-Time Methods in Finance: A Review and an Assessment, in The Journal of Finance, Volume 55, Issue 4, august 2000, pp. 1569-(5)22, online at <http://www.kelley.iu.edu/clundbla/sund.pdf>.
- [Socaciu2009a] Tiberiu Socaciu, Paralelizarea ecuațiilor de tip Black-Scholes si Garman (Parallelization of Black-Scholes and Garman equations, in Romanian), Editura InfoData, will be appear.
- [Socaciu2009b] Tiberiu Socaciu, Ioan Maxim, Algorithm for a Message-based Architecture for Black-Scholes Equation, in IECS 2009, (5)th International Economic Conference "Industrial Revolutions, from the Globalization

634 See [Socaciu2009a], [Chen1996].

635 See [Socaciu2009a], [Iftimie2008].

and Post-Globalization Perspective", 7-8 mai 2009, Sibiu, Romania, section 5, 2009, pp. 236-241, ISBN 978-973-739-775-1.

13. [Socaciu2009c] Tiberiu Socaciu, O rezolvare paralela a ecuatiei Black-Scholes, at Zilele Academice Aradene, 8-10 mai 2009.